



TITLE:

Weak and Strong Convergence Theorems for Normally Generalized Hybrid Mappings in Hilbert Spaces (Study on Nonlinear Analysis and Convex Analysis)

AUTHOR(S):

Takahashi, Wataru

CITATION:

Takahashi, Wataru. Weak and Strong Convergence Theorems for Normally Generalized Hybrid Mappings in Hilbert Spaces (Study on Nonlinear Analysis and Convex Analysis). 数理解析研究所講究録 2019, 2112: 65-72

ISSUE DATE:

2019-04

URL:

<http://hdl.handle.net/2433/251990>

RIGHT:

Weak and Strong Convergence Theorems for Normally Generalized Hybrid Mappings in Hilbert Spaces

慶応義塾大学自然科学研究教育センター, 高雄医科大学基礎科学センター
高橋渉 (Wataru Takahashi)

Keio Research and Education Center for Natural Sciences, Keio University, Japan and
Center for Fundamental Science, Kaohsiung Medical University, Kaohsiung 80702, Taiwan
Email: wataru@is.titech.ac.jp; wataru@a00.itscom.net

Abstract. In this article, using Mann's type iteration, Halpern's type iteration, hybrid method and shrinking projection method, we obtain weak and strong convergence theorems for two generalized hybrid mappings and two normally 2-generalized hybrid mappings in a Hilbert space without assuming that they are commutative.

2010 *Mathematics Subject Classification:* 47H05, 47H09

Keywords and phrases: Fixed point, attractive point, Mann iteration, Halpern iteration, hybrid method, shrinking projection method, generalized hybrid mapping.

1 Introduction

Let H be a real Hilbert space and let C be a nonempty subset of H . A mapping T from C into H is called *generic generalized hybrid* [21] if there exist $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ such that $\alpha + \beta + \gamma + \delta \geq 0$, $\alpha + \beta > 0$ and

$$\alpha \|Tx - Ty\|^2 + \beta \|x - Ty\|^2 + \gamma \|Tx - y\|^2 + \delta \|x - y\|^2 \leq 0$$

for all $x, y \in C$. The class of generic generalized hybrid mappings covers generalized hybrid mappings defined by Kocourek, Takahashi and Yao [6]. A mapping $T : C \rightarrow H$ is called *generalized hybrid* [6] if there exist $\alpha, \beta \in \mathbb{R}$ such that

$$\alpha \|Tx - Ty\|^2 + (1 - \alpha) \|x - Ty\|^2 \leq \beta \|Tx - y\|^2 + (1 - \beta) \|x - y\|^2$$

for all $x, y \in C$. The generalized hybrid mappings were extended by Maruyama, Takahashi and Yao [11] as follows: A mapping $T : C \rightarrow C$ is called *2-generalized hybrid* [11] if there exist $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{R}$ such that

$$\begin{aligned} \alpha_2 \|T^2x - Ty\|^2 + \alpha_1 \|Tx - Ty\|^2 + (1 - \alpha_1 - \alpha_2) \|x - Ty\|^2 \\ \leq \beta_2 \|T^2x - y\|^2 + \beta_1 \|Tx - y\|^2 + (1 - \beta_1 - \beta_2) \|x - y\|^2 \end{aligned}$$

for all $x, y \in C$. Very recently, 2-generalized hybrid mappings were extended by Kondo and Takahashi [7]. A mapping $T : C \rightarrow C$ is called *normally 2-generalized hybrid* [7] if there exist $\alpha_0, \beta_0, \alpha_1, \beta_1, \alpha_2, \beta_2 \in \mathbb{R}$ such that

$$\begin{aligned} & \alpha_2 \|T^2x - Ty\|^2 + \alpha_1 \|Tx - Ty\|^2 + \alpha_0 \|x - Ty\|^2 \\ & + \beta_2 \|T^2x - y\|^2 + \beta_1 \|Tx - y\|^2 + \beta_0 \|x - y\|^2 \leq 0 \end{aligned}$$

for all $x, y \in C$, where $\sum_{n=0}^2 (\alpha_n + \beta_n) \geq 0$ and $\alpha_2 + \alpha_1 + \alpha_0 > 0$.

In this article, using Mann's type iteration, Halpern's type iteration, hybrid method and shrinking projection method, we obtain weak and strong convergence theorems for two generalized hybrid mappings and two normally 2-generalized hybrid mappings in a Hilbert space without assuming that they are commutative.

2 Preliminaries

Throughout this paper, we denote by \mathbb{N} the set of positive integers and by \mathbb{R} the set of real numbers. Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$. We denote the strong convergence and the weak convergence of $\{x_n\}$ to $x \in H$ by $x_n \rightarrow x$ and $x_n \rightharpoonup x$, respectively. In a Hilbert space, it is known that

$$2\langle x - y, y \rangle \leq \|x\|^2 - \|y\|^2 \leq 2\langle x - y, x \rangle \quad (2.1)$$

for all $x, y \in H$ and

$$\|\alpha x + (1 - \alpha)y\|^2 = \alpha \|x\|^2 + (1 - \alpha) \|y\|^2 - \alpha(1 - \alpha) \|x - y\|^2 \quad (2.2)$$

for all $x, y \in H$ and $\alpha \in \mathbb{R}$; see [15]. Furthermore, we have that

$$2\langle x - y, z - w \rangle = \|x - w\|^2 + \|y - z\|^2 - \|x - z\|^2 - \|y - w\|^2 \quad (2.3)$$

for all $x, y, z, w \in H$. Let H be a Hilbert space and let C be a nonempty subset of H . Let T be a mapping of C into H . We denote by $A(T)$ the set of *attractive points* [17] of T , i.e.,

$$A(T) = \{z \in H : \|Tx - z\| \leq \|x - z\|, \quad \forall x \in C\}.$$

We also denote by $F(T)$ the set of fixed points of T . A mapping $T : C \rightarrow H$ with $F(T) \neq \emptyset$ is called *quasi-nonexpansive* if

$$\|Tx - u\| \leq \|x - u\|, \quad \forall x \in C, \quad u \in F(T).$$

If C is closed and convex and $T : C \rightarrow H$ with $F(T) \neq \emptyset$ is quasi-nonexpansive, then $F(T)$ is closed and convex; see Itoh and Takahashi [5]. For a nonempty, closed and convex subset D of H , the nearest point projection of H onto D is denoted by P_D , that is, $\|x - P_Dx\| \leq \|x - y\|$ for all $x \in H$ and $y \in D$. Such a mapping P_D is called the metric projection of H onto D . We know that the metric projection P_D is firmly nonexpansive, i.e.,

$$\|P_Dx - P_Dy\|^2 \leq \langle P_Dx - P_Dy, x - y \rangle$$

for all $x, y \in H$. Furthermore, $\langle x - P_D x, y - P_D x \rangle \leq 0$ holds for all $x \in H$ and $y \in D$; see [14, 15]. Using this inequality and (2.3), we have that

$$\|P_D x - y\|^2 + \|P_D x - x\|^2 \leq \|x - y\|^2, \quad \forall x \in H, y \in D. \quad (2.4)$$

The following result was proved by Takahashi and Toyoda [19].

Lemma 2.1 ([19]). *Let H be a Hilbert space and let C be a nonempty, closed and convex subset of H . Let $\{x_n\}$ be a sequence in H . If $\|x_{n+1} - u\| \leq \|x_n - u\|$ for all $n \in \mathbb{N}$ and $u \in C$, then $\{P_C x_n\}$ converges strongly to $z \in C$, where P_C is the metric projection of H onto C .*

To prove one of our main results, we also need the following lemmas by Aoyama, Kimura, Takahashi and Toyoda [1, 24] and Maingé [9].

Lemma 2.2 ([1, 24]). *Let $\{s_n\}$ be a sequence of nonnegative real numbers, let $\{\alpha_n\}$ be a sequence of $[0, 1]$ with $\sum_{n=1}^{\infty} \alpha_n = \infty$, let $\{\beta_n\}$ be a sequence of nonnegative real numbers with $\sum_{n=1}^{\infty} \beta_n < \infty$, and let $\{\gamma_n\}$ be a sequence of real numbers with $\limsup_{n \rightarrow \infty} \gamma_n \leq 0$. Suppose that*

$$s_{n+1} \leq (1 - \alpha_n)s_n + \alpha_n \gamma_n + \beta_n$$

for all $n = 1, 2, \dots$. Then $\lim_{n \rightarrow \infty} s_n = 0$.

Lemma 2.3 ([9]). *Let $\{X_n\}$ be a sequence of real numbers. Assume that $\{X_n\}$ is not monotone decreasing for sufficiently large $n \in \mathbb{N}$, in other words, there exists a subsequence $\{X_{n_i}\}$ of $\{X_n\}$ such that $X_{n_i} < X_{n_i+1}$ for all $i \in \mathbb{N}$. Let $n_0 \in \mathbb{N}$ such that $\{k \leq n_0 : X_k < X_{k+1}\} \neq \emptyset$. Define a sequence $\{\tau(n)\}_{n \geq n_0}$ of natural numbers as follows:*

$$\tau(n) = \max \{k \leq n : X_k < X_{k+1}\}, \quad \forall n \geq n_0.$$

Then, the followings hold:

- (i) $\tau(n) \rightarrow \infty$ as $n \rightarrow \infty$;
- (ii) $X_n \leq X_{\tau(n)+1}$ and $X_{\tau(n)} < X_{\tau(n)+1}$, $\forall n \geq n_0$.

3 Weak convergence theorems of Mann's type iteration

In this section, using Lemma 2.1, we obtain a weak convergence theorem of Mann's type iteration [10] for finding a common attractive point of two generalized hybrid mappings without assuming that the mappings are commutative. Before proving the theorem, we need the following lemma.

Lemma 3.1. *Let H be a Hilbert space and let C be a nonempty subset of H . Let $T : C \rightarrow H$ be a generalized hybrid mapping and let $\{x_n\} \subset C$. If $x_n \rightarrow z$ and $x_n - Tx_n \rightarrow 0$, then $z \in A(T)$. Additionally, if C is closed and convex, then $z \in F(T)$.*

Theorem 3.2 ([16]). *Let H be a Hilbert space and let C be a nonempty and convex subset of H . Let S and T be generalized hybrid mappings of C into itself such that $A(S) \cap A(T) \neq \emptyset$. Let $\{x_n\}$ be a sequence generated by $x_1 = x \in C$ and*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n)(\gamma_n Sx_n + (1 - \gamma_n)Tx_n), \quad \forall n \in \mathbb{N},$$

where $a, b, c, d \in \mathbb{R}$, $\{\gamma_n\}$ and $\{\alpha_n\}$ satisfy the following:

$$0 < a \leq \gamma_n \leq b < 1 \quad \text{and} \quad 0 < c \leq \alpha_n \leq d < 1, \quad \forall n \in \mathbb{N}.$$

Then $\{x_n\}$ converges weakly to a point $z \in A(S) \cap A(T)$, where $z = \lim_{n \rightarrow \infty} P_{A(S) \cap A(T)} x_n$. Additionally, if C is closed, then $\{x_n\}$ converges weakly to a point $z \in F(S) \cap F(T)$, where $z = \lim_{n \rightarrow \infty} P_{F(S) \cap F(T)} x_n$.

We can also prove a weak convergence theorem by Mann's type iteration [10] for noncommutative two normally 2-generalized hybrid mappings in Hilbert spaces; see also [3].

Theorem 3.3 ([13]). *Let H be a Hilbert space and let C be a nonempty and convex subset of H . Let S and T be normally 2-generalized hybrid mappings of C into itself such that $A(S) \cap A(T) \neq \emptyset$. Given $x_1 \in C$, define a sequence $\{x_n\}$ in C as follows:*

$$x_{n+1} = a_n x_n + b_n (\gamma_n S + (1 - \gamma_n) T) x_n + c_n (\delta_n S^2 + (1 - \delta_n) T^2) x_n$$

for all $n \in \mathbb{N}$, where $a, b, c, d, e, f \in \mathbb{R}$ and $\{\gamma_n\}, \{\delta_n\}, \{a_n\}, \{b_n\}, \{c_n\} \subset [0, 1]$ satisfy the following:

$$0 < a \leq \gamma_n \leq b < 1, \quad 0 < c \leq \delta_n \leq d < 1, \\ a_n + b_n + c_n = 1 \quad \text{and} \quad 0 < e \leq a_n, b_n, c_n \leq f < 1, \quad \forall n \in \mathbb{N}.$$

Then $\{x_n\}$ converges weakly to a point u of $A(S) \cap A(T)$, where $u = \lim_{n \rightarrow \infty} P_{A(S) \cap A(T)} x_n$.

Additionally, if C is closed, then $\{x_n\}$ converges weakly to a point $z \in F(S) \cap F(T)$, where $z = \lim_{n \rightarrow \infty} P_{F(S) \cap F(T)} x_n$.

4 Strong convergence theorems of Halpern's type iteration

In this section, using Lemmas 2.2 and 2.3, we prove the following strong convergence theorem of Halpern's type iteration [2] for noncommutative two generalized hybrid mappings in a Hilbert space; see also [22].

Theorem 4.1 ([16]). *Let H be a Hilbert space and let C be a nonempty and convex subset of H . Let S and T be generalized hybrid mappings of C into itself with $A(S) \cap A(T) \neq \emptyset$. Given $x_1 \in C$ and $\{u_n\} \subset C$ with $u_n \rightarrow u$, define a sequence $\{x_n\}$ in C as follows:*

$$x_{n+1} = \alpha_n u_n + (1 - \alpha_n) (\beta_n x_n + (1 - \beta_n) (\gamma_n S x_n + (1 - \gamma_n) T x_n))$$

for all $n \in \mathbb{N}$, where $a, b, c, d \in \mathbb{R}$, $\{\gamma_n\}$, $\{\alpha_n\}$ and $\{\beta_n\}$ satisfy the following:

$$\lim_{n \rightarrow \infty} \alpha_n = 0, \quad \sum_{n=1}^{\infty} \alpha_n = \infty,$$

$$0 < a \leq \gamma_n \leq b < 1 \quad \text{and} \quad 0 < c \leq \beta_n \leq d < 1, \quad \forall n \in \mathbb{N}.$$

Then the sequence $\{x_n\}$ converges strongly to $P_{A(S) \cap A(T)} u$, where $P_{A(S) \cap A(T)}$ is the metric projection from H onto $A(S) \cap A(T)$. Additionally, if C is closed, then $\{x_n\}$ converges strongly to $P_{F(S) \cap F(T)} u$, where $P_{F(S) \cap F(T)}$ is the metric projection from H onto $F(S) \cap F(T)$.

We can also prove a strong convergence theorem by Halpern's type iteration [2, 23] for noncommutative two normally 2-generalized hybrid mappings in Hilbert spaces; see also [3, 8].

Theorem 4.2 ([13]). *Let H be a Hilbert space and let C be a nonempty and convex subset of H . Let S and T be normally 2-generalized hybrid mappings of C into itself such that $A(S) \cap A(T) \neq \emptyset$. Given $x_1, z \in C$, define a sequence $\{x_n\}$ in C as follows:*

$$\begin{cases} x_{n+1} = \lambda_n z + (1 - \lambda_n) z_n, \\ z_n = a_n x_n + b_n (\gamma_n S + (1 - \gamma_n) T) x_n + c_n (\delta_n S^2 + (1 - \delta_n) T^2) x_n, \end{cases} \quad \forall n \in \mathbb{N},$$

where $a, b, c, d, e, f \in \mathbb{R}$ and $\{\lambda_n\}, \{\gamma_n\}, \{\delta_n\}, \{a_n\}, \{b_n\}, \{c_n\} \subset [0, 1]$ satisfy the following:

$$\begin{aligned} \lim_{n \rightarrow \infty} \lambda_n &= 0, \quad \sum_{n=1}^{\infty} \lambda_n = \infty, \\ 0 &< a \leq \gamma_n \leq b < 1, \quad 0 < c \leq \delta_n \leq d < 1, \\ a_n + b_n + c_n &= 1 \quad \text{and} \quad 0 < e \leq a_n, b_n, c_n \leq f < 1, \quad \forall n \in \mathbb{N}. \end{aligned}$$

Then the sequence $\{x_n\}$ converges strongly to $z_0 = P_{A(S) \cap A(T)} z$, where $P_{A(S) \cap A(T)}$ is the metric projection from H onto $A(S) \cap A(T)$.

Additionally, if C is closed, then $\{x_n\}$ converges strongly to $P_{F(S) \cap F(T)} z$, where $P_{F(S) \cap F(T)}$ is the metric projection from H onto $F(S) \cap F(T)$.

5 Strong convergence theorems by hybrid methods

In this section, we obtain a strong convergence theorem by the hybrid method [12] for finding a common fixed point of two generalized hybrid mappings without assuming that the mappings are commutative.

Theorem 5.1 ([4]). *Let H be a Hilbert space and let C be a nonempty, closed and convex subset of H . Let $S, T : C \rightarrow C$ be generalized hybrid mappings such that $F(S) \cap F(T) \neq \emptyset$. Let $\{x_n\} \subset C$ be a sequence generated by $x_1 \in C$ and*

$$\begin{cases} y_n = \alpha_n x_n + (1 - \alpha_n) (\gamma_n S x_n + (1 - \gamma_n) T x_n), \\ C_n = \{z \in C : \|y_n - z\| \leq \|x_n - z\|\}, \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\}, \\ x_{n+1} = P_{C_n \cap Q_n} x_1, \end{cases} \quad \forall n \in \mathbb{N},$$

where $P_{C_n \cap Q_n}$ is the metric projection of H onto $C_n \cap Q_n$ and $a, b, c \in \mathbb{R}$ and $\{\alpha_n\}, \{\gamma_n\} \subset [0, 1]$ satisfy

$$0 \leq \alpha_n \leq a < 1 \quad \text{and} \quad 0 < b \leq \gamma_n \leq c < 1, \quad \forall n \in \mathbb{N}.$$

Then $\{x_n\}$ converges strongly to $z_0 = P_{F(S) \cap F(T)} x_1$, where $P_{F(S) \cap F(T)}$ is the metric projection of H onto $F(S) \cap F(T)$.

Next, we prove a strong convergence theorem by the shrinking projection method [18] for noncommutative two generalized hybrid mappings in a Hilbert space.

Theorem 5.2 ([4]). *Let H be a Hilbert space and let C be a nonempty, closed and convex subset of H . Let $S, T : C \rightarrow C$ be generalized hybrid mappings such that $F(S) \cap F(T) \neq \emptyset$. Let $\{u_n\}$ be a sequence in C such that $u_n \rightarrow u$. Let $C_1 = C$ and let $\{x_n\} \subset C$ be a sequence generated by $x_1 \in C$ and*

$$\begin{cases} y_n = \alpha_n x_n + (1 - \alpha_n)(\gamma_n Sx_n + (1 - \gamma_n)Tx_n), \\ C_{n+1} = \{z \in C_n : \|y_n - z\| \leq \|x_n - z\|\}, \\ x_{n+1} = P_{C_{n+1}} u_{n+1}, \quad \forall n \in \mathbb{N}, \end{cases}$$

where $P_{C_{n+1}}$ is the metric projection of H onto C_{n+1} and $b, c \in \mathbb{R}$ and $\{\alpha_n\}, \{\gamma_n\} \subset [0, 1]$ satisfy

$$0 \leq \liminf_{n \rightarrow \infty} \alpha_n < 1 \quad \text{and} \quad 0 < b \leq \gamma_n \leq c < 1, \quad \forall n \in \mathbb{N}.$$

Then, $\{x_n\}$ converges strongly to $z_0 = P_{F(S) \cap F(T)} u$, where $P_{F(S) \cap F(T)}$ is the metric projection of H onto $F(S) \cap F(T)$.

Furthermore, using the hybrid method [12], we prove a strong convergence theorem for noncommutative normally 2-generalized hybrid mappings in a Hilbert space.

Theorem 5.3 ([20]). *Let H be a Hilbert space and let C be a nonempty, closed and convex subset of H . Let $S, T : C \rightarrow C$ be normally 2-generalized hybrid mappings such that $F(S) \cap F(T) \neq \emptyset$. Let $\{x_n\} \subset C$ be a sequence generated by $x_1 \in C$ and*

$$\begin{cases} y_n = a_n x_n + b_n(\gamma_n S + (1 - \gamma_n)T)x_n + c_n(\delta_n S^2 + (1 - \delta_n)T^2)x_n, \\ C_n = \{z \in C : \|y_n - z\| \leq \|x_n - z\|\}, \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\}, \\ x_{n+1} = P_{C_n \cap Q_n} x_1, \quad \forall n \in \mathbb{N}, \end{cases}$$

where $P_{C_n \cap Q_n}$ is the metric projection of H onto $C_n \cap Q_n$ and $a, b, c, d, e, f \in \mathbb{R}$ and $\{\gamma_n\}, \{\delta_n\}, \{a_n\}, \{b_n\}, \{c_n\} \subset [0, 1]$ satisfy the following:

$$0 < a \leq \gamma_n \leq b < 1, \quad 0 < c \leq \delta_n \leq d < 1, \\ a_n + b_n + c_n = 1 \quad \text{and} \quad 0 < e \leq a_n, b_n, c_n \leq f < 1, \quad \forall n \in \mathbb{N}.$$

Then $\{x_n\}$ converges strongly to $z_0 = P_{F(S) \cap F(T)} x_1$, where $P_{F(S) \cap F(T)}$ is the metric projection of H onto $F(S) \cap F(T)$.

Finally, we prove a strong convergence theorem by the shrinking projection method [18] for noncommutative normally 2-generalized hybrid mappings in a Hilbert space.

Theorem 5.4 ([20]). *Let H be a Hilbert space and let C be a nonempty, closed and convex subset of H . Let $S, T : C \rightarrow C$ be normally 2-generalized hybrid mappings such that $F(S) \cap F(T) \neq \emptyset$. Let $\{u_n\}$ be a sequence in C such that $u_n \rightarrow u$. Let $C_1 = C$ and let $\{x_n\} \subset C$ be a sequence generated by $x_1 \in C$ and*

$$\begin{cases} y_n = a_n x_n + b_n(\gamma_n S + (1 - \gamma_n)T)x_n + c_n(\delta_n S^2 + (1 - \delta_n)T^2)x_n, \\ C_{n+1} = \{z \in C_n : \|y_n - z\| \leq \|x_n - z\|\}, \\ x_{n+1} = P_{C_{n+1}} u_{n+1}, \quad \forall n \in \mathbb{N}, \end{cases}$$

where $P_{C_{n+1}}$ is the metric projection of H onto C_{n+1} and $a, b, c, d, e, f \in \mathbb{R}$ and $\{\gamma_n\}, \{\delta_n\}, \{a_n\}, \{b_n\}, \{c_n\} \subset [0, 1]$ satisfy the following:

$$0 < a \leq \gamma_n \leq b < 1, \quad 0 < c \leq \delta_n \leq d < 1, \\ a_n + b_n + c_n = 1 \quad \text{and} \quad 0 < e \leq a_n, b_n, c_n \leq f < 1, \quad \forall n \in \mathbb{N}.$$

Then $\{x_n\}$ converges strongly to $z_0 = P_{F(S) \cap F(T)}u$, where $P_{F(S) \cap F(T)}$ is the metric projection of H onto $F(S) \cap F(T)$.

Acknowledgements. The author was partially supported by Grant-in-Aid for Scientific Research No. 15K04906 from Japan Society for the Promotion of Science.

References

- [1] K. Aoyama, Y. Kimura, W. Takahashi and M. Toyoda, *Approximation of common fixed points of a countable family of nonexpansive mappings in a Banach space*, Nonlinear Anal. **67** (2007), 2350-2360.
- [2] B. Halpern, *Fixed points of nonexpanding maps*, Bull. Amer. Math. Soc. **73** (1967), 957-961.
- [3] M. Hojo, A. Kondo and W. Takahashi, *Weak and strong convergence theorems for commutative normally 2-generalized hybrid mappings in Hilbert spaces*, Linear Nonlinear Anal. **4** (2018), 145-156.
- [4] M. Hojo and W. Takahashi, *Strong convergence theorems by hybrid methods for noncommutative two nonlinear mappings in Hilbert spaces*, in Nonlinear Analysis and Convex Analysis, Yokohama Publishers, Yokohama, to appear.
- [5] S. Itoh and W. Takahashi, *The common fixed point theory of singlevalued mappings and multivalued mappings*, Pacific J. Math. **79** (1978), 493-508.
- [6] P. Kocourek, W. Takahashi and J.-C. Yao, *Fixed point theorems and weak convergence theorems for generalized hybrid mappings in Hilbert spaces*, Taiwanese J. Math. **14** (2010), 2497-2511.
- [7] A. Kondo and W. Takahashi, *Attractive point and weak convergence theorems for normally N-generalized hybrid mappings in Hilbert spaces*, Linear Nonlinear Anal. **3** (2017), 297-310.
- [8] A. Kondo and W. Takahashi, *Strong convergence theorems of Halpern's type for normally 2-generalized hybrid mappings in Hilbert spaces*, J. Nonlinear Convex Anal. **19** (2018), 617-631.
- [9] P. E. Maingé, *Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization*, Set-Valued Anal. **16** (2008), 899-912.
- [10] W. R. Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. **4** (1953), 506-510.
- [11] T. Maruyama, W. Takahashi and M. Yao, *Fixed point and mean ergodic theorems for new nonlinear mappings in Hilbert spaces*, J. Nonlinear Convex Anal. **12** (2011), 185-197.
- [12] K. Nakajo and W. Takahashi, *Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups*, J. Math. Anal. Appl. **279** (2003), 372-378.
- [13] S. Takahashi and W. Takahashi, *Weak and strong convergence theorems for noncommutative normally 2-generalized hybrid mappings in Hilbert spaces*, J. Nonlinear Convex Anal. **19** (2018), 1427-1441.

- [14] W. Takahashi, *Nonlinear Functional Analysis*, Yokohama Publishes, Yokohama, 2000.
- [15] W. Takahashi, *Introduction to Nonlinear and Convex Analysis*, Yokohama Publishes, Yokohama, 2009.
- [16] W. Takahashi, *Weak and strong convergence theorems for noncommutative two generalized hybrid mappings in Hilbert spaces*, J. Nonlinear Convex Anal. **19** (2018), 867-880.
- [17] W. Takahashi and Y. Takeuchi, *Nonlinear ergodic theorem without convexity for generalized hybrid mappings in a Hilbert space*, J. Nonlinear Convex Anal. **12** (2011), 399-406.
- [18] W. Takahashi, Y. Takeuchi and R. Kubota, *Strong convergence theorems by hybrid methods for families of nonexpansive mappings in Hilbert spaces*, J. Math. Anal. Appl. **341** (2008), 276-286.
- [19] W. Takahashi and M. Toyoda, *Weak convergence theorems for nonexpansive mappings and monotone mappings*, J. Optim. Theory Appl. **118** (2003), 417-428.
- [20] W. Takahashi, C.-F. Wen and J.-C. Yao, *Strong convergence theorems by hybrid methods for noncommutative normally 2-generalized hybrid mappings in Hilbert spaces*, Appl. Anal. Optim., to appear.
- [21] W. Takahashi, N. C. Wong and J.-C. Yao, *Attractive point and weak convergence theorems for new generalized hybrid mappings in Hilbert spaces*, J. Nonlinear Convex Anal. **13** (2012), 745-757.
- [22] W. Takahashi, N. C. Wong and J.-C. Yao, *Attractive points and Halpern-type strong convergence theorems in Hilbert spaces*, J. Fixed Point Theory Appl. **17** (2015), 301-311.
- [23] R. Wittmann, *Approximation of fixed points of nonexpansive mappings*, Arch. Math. **58** (1992), 486-491.
- [24] H. K. Xu, *Another control condition in an iterative method for nonexpansive mappings*, Bull. Aust. Math. Soc. **65** (2002), 109-113.